

reddish ochraceous, the last with the disk paler, and with the anterior half of lateral margin and an apical spot levigate pale ochraceous; eyes, apex of second joint of antennæ, and third and fourth joints (excluding bascs) black; cuneus reddish; membrano pale brownish ochraceous and semihyaline, the cellular margins brownish; head beneath and sternum reddish ochraceous; abdomen, metasternum, and centre of mesosternum (*carded specimen*) black; legs ochraceous.

Long.  $3\frac{1}{2}$  millim.

*Hab.* GUATEMALA, Las Mercedes (*Champion*).

b. *Posterior margin of the pronotum truncate.*

4. *Carnus funebris*, n. sp.

Bright greenish black; basal half of head, antennæ, rostrum, and legs ochraceous; rostrum with the apex piceous; antennæ with the apex of the second joint luteous (third and fourth joints mutilated); pronotum very coarsely punctate; membrane fuscous, cellular margins pale brownish; abdomen beneath having the transverse segmental margins somewhat paler.

Long. 4 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

PSEUDOCARNUS, gen. nov.

Differing from *Carnus* by having the basal joint of the antennæ longer than the head and incrassated; second joint about twice as long as the first; third and fourth joints very short, and together scarcely or about as long only as the first joint. Pronotum obscurely punctate.

1. *Pseudocarnus lineolatus*, n. sp. (Tab. XXVIII. fig. 8.)

Ochraceous; antennæ with the basal joint reddish; apical fourth of second joint, the third joint and fourth joint (excluding base and apex) castaneous; eyes silvery grey; pronotum with the posterior margin pale sanguineous, the lateral posterior margins black; inner (excluding base) and sutural margins of clavus, a basal streak and submarginal fascia (deflected at centre and angularly continued to near claval apex) to corium black; cuneus with the lateral margins and apex sanguineous; membrane pale fuscous and semihyaline, the cellular margins black. Body beneath and legs ochraceous, basal lateral margins of prosternum black.

Long. 4 millim.

*Hab.* PANAMA, Bugaba (*Champion*).

This species is peculiar by the great length of the first and second joints of the antennæ and the relative shortness of the third and fourth, which together are not much more than half the length of the second.

2. *Pseudocarnus magnus*, n. sp. (Tab. XXVIII. fig. 9.)

Ochraceous; eyes, lateral margins of pronotum, a narrow central linear fascia to clavus, a sublateral fascia to corium deflected a short distance from apex and transversely continued to inner apical angle, castaneous; antennæ ochraceous, basal joint and apex of second joint pale castaneous, third joint ochraceous, fourth fuscous; clavus and inner disk of corium much suffused with reddish; cuneus with the inner margin sanguineous; membrane pale brownish and semihyaline, cellular margins castaneous; sternum and legs ochraceous; abdomen reddish.

Long. 6 millim.

*Hab.* GUATEMALA, Cerro Zunil (*Champion*).