

(excepting basal angular spots and lateral margins and apex), margins of clavus, a reticulated transverse fascia to corium, a transverse streak at base of cuneus, cellular margins to membrane, sternal and abdominal sutures (*carded specimen*) black; anterior and intermediate legs ochraceous (posterior legs mutilated).

Long. $5\frac{1}{2}$ millim.

Hab. GUATEMALA, Sinanja (*Champion*).

14. *Neofurius pallidulus*, n. sp. (Tab. XXVIII. fig. 20.)

Creamy white; eyes, apex of second joint, apical half of third joint and the fourth joint of antennæ, inner margins of clavus, and an oblique spot near centre of costal margin of corium, black; costal margin of corium from near base to centre, and posterior femora dark sanguineous; lateral margins of cuneus narrowly fuscous; body beneath indistinctly seen (*carded specimen*).

Costal margin of corium amplified and possessing a distinct pouch-like depression about centre.

Long. 4 millim.

Hab. GUATEMALA, Panajachel (*Champion*).

15. *Neofurius plagosus*, n. sp. (Tab. XXVIII. fig. 21.)

Creamy white, tinged with very pale ochraceous; eyes, first and second joints of antennæ, a rounded spot at base, and an elongate spot about centre of clavus, an oblique fascia about centre of corium, inner margin of cuneus, cellular margins to membrane, two lateral spots to sternum, apices of femora, and the tibiæ fuscous or black; base of first joint of antennæ and apex of posterior tibiæ creamy white, tinged with pale ochraceous.

Long. 6 millim.

Hab. PANAMA, Volcan de Chiriqui 2500 to 4000 feet (*Champion*).

16. *Neofurius linearis*, n. sp. (Tab. XXVIII. fig. 22.)

Head, pronotum, and legs ochraceous; scutellum, corium, and membrane creamy white; eyes, first and second joints of antennæ, a longitudinal fascia crossing clavus and continued along corium to inner angle of cuneus, outer and inner margins of cuneus, costal margin of corium, cellular margins to membrane, apices of femora, the tibiæ, and apices of tarsi fuscous or black; base of first joint of antennæ pale ochraceous; body beneath pale ochraceous or with the abdomen sometimes creamy white.

Long. 5 millim.

Hab. GUATEMALA, San Gerónimo, Mirandilla (*Champion*).

BIBACULUS, gen. nov.

Allied to *Neofurius*, but with the body elongate and very indistinctly pilose, the lateral margins of the corium parallel, not dilated. Antennæ with the first joint about as long as or a little shorter than the head, second joint almost twice as long as the first, third joint distinctly longer than second. Pronotum as in preceding species, but longer, posterior margin concavely truncate, and about twice the width of anterior margin. Other characters as in *Neofurius*.

This genus has apparently considerable superficial resemblance to *Hyaliodes*, Reut., founded for the reception of *Capsus vitripennis*, Say, a species I only know by the figure given by Townend Glover (Ill. Ins. Hem. t. vii. f. 8). Reuter's genus is, however, rendered very distinct by the great length of the basal joint of the antennæ.