

*Plagiotylus* Sc.

*P. dispar* Rt. – Palestine (Bodenheimer op.cit.); Jericho, Transjordanian, some spec., J. Sahlberg (!). – Endemic.

*Dimorphocoris* Rt.

*D. punctiger* (Hv.) – Palestine (BODENHEIMER op.cit.). – Endemic.

*D. lateralis* Rt. – Haifa, 1 spec., J. Sahlberg (!). Pontomediterranean. Known from Crete and Israel.

The species resembles *D. punctiger*, but the antennae are shorter, the proportions between the joints being 14 + 29 + 20 + 10 (♂) (in *punctiger* ♂ 19 + 40 + 29 + 15, 1 unit = 0.088 mm.). The tibial spines arise from small dark spots in *D. lateralis* also. I have not seen the type specimen from Crete. The specimen from Haifa has been illustrated by LINDBERG (1956, p. 59).

*D. mariae* Lv. – Galilea, Vall. Kison, 3 spec., 31. III. 1904, J. Sahlberg (!); Palestine, 2 spec., Bodenheimer (!). – Endemic.

The male much resembles the corresponding sex of *D. debilis* (Rt.) but is bigger, length 4.5 – 4.6 mm. (*debilis* ♂ 3.9 – 4.2 mm.) and the 2nd and 3rd antennal joints are longer, the proportions between the joints being 13 + 35 + 25 + 9 (in *debilis* ♂ 12 + 30 + 20 + 8, 1 unit = 0.088 mm.).

*D. debilis* (Rt.) – Palestine (BODENHEIMER op.cit.); Haifa, 1 spec., J. Sahlberg (!); Ness Zionah, 2 spec., 15. II – 3. III. 1933, Carmin (!); Palestine, 7 spec., Bodenheimer (!). – Holomediterranean.

*Orthocephalus* Fb.

*O. proserpinae* Ms. R. – Palestine (BODENHEIMER op.cit.). – Holomediterranean.

*O. tenuicornis* (Ms. R.) – Palestine (BODENHEIMER op.cit.); Beersheba, 1 spec., Bytinski-Salz (!); Galilea, some spec., J. Sahlberg (!); Jerusalem, 2 spec., 23. IV. 1946, Houška (HOBERLANDT 1951, p. 32); Kiriath Anavim, 2 spec., 3. VI. 1931, Bodenheimer (!); Miqve Israel, 2 spec., Bodenheimer (!); Ramath Gan, 15 spec., 22. III. 1942, Houška (HOBERLANDT op.cit.). – Holomediterranean.

*Pachytomella* Rt.

*P. phoenicea* (Hv.) – Palestine (BODENHEIMER op.cit.); El Hamme, 1 spec., 21. III. 1945, Bytinski-Salz (!); Galilea, some spec., Saalas (!). – Syrio-Anatolian.

*P. passerinii* (C.) – Palestine (BODENHEIMER op.cit.). – Holomediterranean.

*Strongylocoris* Blanch.

*S. niger* (H. S.) – Palestine (BODENHEIMER op.cit.). – Holomediterranean, extending far into Central Europe.

*S. amabilis* (Dgl. Sc.) – Palestine (BODENHEIMER op.cit.); Benjamina, 1 spec., 23. III. 1942, Houška (HOBERLANDT 1951, p. 31). – Syrio-Anatolian.

*S. cicadifrons* C. – Palestine (BODENHEIMER op.cit.); Palestine, 3 spec., Bodenheimer (!). – Holomediterranean.

*Halticus* Hhn.

*H. rugosus* Rt. – Palestine (BODENHEIMER op.cit.). – Endemic.