



Fig. 7. *Amblytylus inscriptus* n.sp.: a head, lateral view; b theca; c left stylus; d vesica. - *A. gregarius* n.sp.: e theca; f claw; g head, lateral view. - Orig.

M. molliculus (Fn.) - Palestine (BODENHEIMER op.cit.); Rehovot, 3 spec., 20. IV. 1958, Swirski (!). - European.

(*M. longirostris* (Fb.)) - Jericho, Transjordanian, 1 spec., J. Sahlberg (!). - Holomediterranean. Not previously recorded from the Middle East.

M. krueperi (Rt.) - Sa'ad, 1 spec., 20. III. 1958, Rubin (!); Jericho, Transjordanian, 1 spec., J. Sahlberg (!). - Holomediterranean. Not previously known from Israel.

Amblytylus Fb.

A. inscriptus n.sp.

♂. Length 3.8 mm. Head yellow-greenish; stylus, median part and margins of frons and base of vertex yellowish. Eyes light grey. Antennae yellowish, 3rd and 4th joints (sometimes also 2nd joint) slightly infuscate. Pronotum yellowish; a median stripe and a transverse stripe behind calli green, forming a cross-shaped green figure on the disk; lateral margins with a green mark, basal lateral angles whitish yellow. Sometimes the calli are slightly infuscate and the greenish area larger, sending 4 broad greenish-yellow branches caudad. Scutellum yellowish with 2 small greenish basal spots; an orange median stripe sometimes present. Clavus, corium and cuneus greenish yellow, lateral margin of corium lighter; an obscure lighter longitudinal stripe present on corium; membrane milky, apical part with a broad bow-shaped smoky area extending to the apices of the membranal cells, which are also distinctly darkened, a dark smoky spot also present in the mediobasal angle of the membrane. Under surface greenish and yellowish. Legs yellowish, femora minutely spotted with dark apically; 1st and 2nd tarsal joints brownish, 3rd joint dark.

A small, slender species. Body elongate, 3.8 × as long as broad. Head 1.3 × as broad as long, in lateral aspect less steeply sloping ventrad apically than in the following species (fig. 7 a);