

pronotum $1.2 \times (\text{♂})$ or $1.4 \times (\text{♀})$ as long as the 2nd antennal joint. *D. pilipes* Rt. has long, erect hairs on the 2nd antennal joint, a black puncturing on the upper surface, etc.

Trigonotylus pallidicornis Rt. — Homs, 2 spec., 27. VIII. 1962. New to Libya. Eremian.

Cyrtopeltis tenuis Rt. — Homs, some, 27. VIII. 1962.

Orthotylus junipericola Lv. ssp. *contractulus* n. ssp.

♂ 2.6 mm, ♀ 2.3 mm. As the nominate form, but vertex $1.9 \times (\text{♂})$ or $2 \times (\text{♀})$ as broad as eye; 2nd antennal joint $0.88 \times$ as long as basal width of pronotum and with dissimilar genitalia: right stylus (fig. 1 h—i) with a basal lobe bearing 3 teeth; hypophysis sharp and directed laterad; the body of the stylus is bluntly produced apical above the hypophysis. Left stylus (fig. 1 j) with hypophysis strongly triangularly produced in upper angle, while the basal angle is roundedly rectangular. Vesical appendages as in the nominate form.

El Merj, 1 ♂ type, 4 paratypes, 31. VIII. 1962. On *Juniperus phoenicea*.

Brachynotocoris cyprius E. WGN. ssp. *inermis* n. ssp.

As nominate form, but styli of ♂ different: Right stylus (fig. 1 k) bearing only one strong tooth, while the apex of the stylus is provided with only a very faint tooth (well developed in the nominate form); the left stylus (fig. 1 l) has a more produced sensory lobe and the upper margin of the stylus (broad aspect) is strongly insinuated.

El Merj—Gubba road, 1 ♂ type, 2 paratypes, 31. VIII. 1962. On *Olea europaea*. The nominate form known from Cyprus and Israel.

Platyseranus putoni Rt. — nr. Tripolis, 1 spec., 27. VIII. 1962; Zouara—Tripolis road, some, 27. VIII. 1962. On *Retama*. New to Libya. Eremian.

Dimorphocoris bergevini LDB. — Djebel Nefoussa, 1 spec., 1. V. 1961, ECKERLEIN. Previously known from Algeria.

Stenoparia putoni FB. — Kussabat, 2 spec., 3. V. 1961; Kseia, 3 spec., 2—5. VI. 1961, ECKERLEIN. Holomediterranean, not previously recorded from Libya.

Roudairea crassicornis Pt. Rt. — Djebel Nefoussa, 1 spec., 1. V. 1961, ECKERLEIN. Eremian, previously known from Algeria and Tunisia.

Megalocoleus cunealis Rt. — Kseia, 1 spec., 2—3. V. 1961, ECKERLEIN. Eremian, previously known from Algeria.

Amblytylus vittiger Rt. — Zouara, many, 29. IV. 1961, ECKERLEIN. Previously known from Algeria.

Tytthus parviceps (Rt.) — El Merj, 1 spec., 30. VIII. 1962. At lamp. New to Libya. Intertropical.

Psallus varians (H. S.) — Wadi-el-Kuf, some, 2. V. 1963 and Gubba, 1 spec., 2. V. 1963, ECKERLEIN. European. New record for Libya.

P. aurora (M. R.) — nr. Benghazi, 1 spec., 30. VIII. 1962. Swept from desert vegetation, probably from *Artemisia* sp. New to Libya. Westmediterranean.

P. (Compsidolon) scutellaris Rt. — Tarhuna, 1 spec., 4. V. 1963, ECKERLEIN. Previously known from Algeria.

P. (Compsidolon) saundersi Rt. — 10 km S of Tripolis, 1 spec., 30. IV. 1961, ECKERLEIN. Previously known from Algeria.

Malacotes phlomidis LDB. — Wadi-el-Kuf, some, 2. V. 1963, ECKERLEIN. On *Phlomis* sp. Westmediterranean. New record for Libya.

Campylomma nicolasi Pt. Rt. — Homs, 1 spec., 27. VIII. 1962; 100 km E of Homs, 2 spec., 28. VIII. 1962.

C. oertzeni Rt. — 100 km E of Homs, 1 spec., 28. VIII. 1962.

Auchenocrepis alboscutellata Pt. — Zouara—Tripolis road, 2 spec., 27. VIII. 1962.