

- A. lineolatus* (Gz.) — Akçakoca road, 1 spec., 12. VIII. 1963; Silivri, 1 spec., 15. VIII. 1963.
- Dichroscytus cyprius* LDB. — Gülek, 1 spec., 6. VIII. 1963. On *Juniperus phoenicea*.
- Lygus pratensis* (L.) — Akçakoca road, some, 12. VIII. 1963.
- L. gemellatus* (H. S.) — Greece, nr. Amfipolis, some, 16. VIII. 1963; Lake Bourou, 2 spec., 16. VIII. 1963.
- L. rugulipennis* POP. — Akçakoca road, 1 spec., 12. VIII. 1963.
- L. pallidulus* (BLANCH.) — nr. Gülek, 2 spec., 6. VIII. 1963.
- Camptozygum aequalis* (VL.) — Yauladagi, 1 spec., 5. VIII. 1963. On *Pinus halepensis*. Not previously recorded from Turkey. European.
- Polymerus cognatus* (FB.) — Ulukişla, 1 spec., 7. VIII. 1963.
- Deracoricus ruber* (L.) — nr. Gerede, 1 spec., 11. VIII. 1963. New to Turkey. Holarctic.
- D. serenus* (DGL. SC.) — 50 km S of Aksaray, some, 7. VIII. 1963; Ulukişla, some, 7. VIII. 1963.
- D. cyprius* E. WGN. — Yauladagi, 3 spec., 5. VIII. 1963. On *Pinus halepensis*. Previously known only from Cyprus.
- D. lutescens* (SCHL.) — nr. Ankara, some, 10. VIII. 1963.
- Alloetomus gothicus* (FN.) — nr. Gerede, some, 11. VIII. 1963. On *Pinus* sp. New to Turkey. European.
- Trigonotylus pulchellus* (H.) — Iskenderun, 2 spec., 6. VIII. 1963. New to Turkey. Ponto-mediterranean with a large extension into Central Europe.
- Fulvius punctum-album* (R.) — Pozanti, 1 spec., 7. VIII. 1963. Under loose bark of a *Pinus* stub.
- Macrolophus caliginosus* E. WGN. — Yauladagi—Belen, some 5. VIII. 1963.
- Cyrtopeltis turcica* SD. — Yauladagi—Belen, 2 spec., 5. VIII. 1963.
- Acrorrhinium conspersum* NH.
- ♀. Fig. 24. 4.2 mm. Dull. Dark coffee-brown. With minute light irroration on head, pronotum and scutellum. 1st antennal joint blackish brown, with a few (3) seta-bearing whitish knobs; 2nd joint greyish brown. Pronotum with a light broken median stripe. Elytra with scanty faint light irroration; a blackish unirrorate triangle starting near the commissural margin of clavus, extending onto corium and continuing as an oblique band near to the costal margin; clavus at tip of scutellum with a roughly \cap -shaped whitish figure bordering the blackish triangle; also corium with a more indistinct lighter area just basad of the blackish figure; apicolateral angle of corium whitish; cuneus blackish brown, somewhat shining; membrane dark smoky, veins concolorous. Legs dark coffee-brown, knees lighter.
- Resembling *Phytocoris* in general appearance. Head (fig 15 j—k) relatively small, $0.7 \times$ as broad as basal width of pronotum. Vertex $1.5 \times$ as broad as eye, slightly convex and with a faint median sulcus. In lateral aspect the head is a little higher than long with the tylus remarkably prominent. Proportions between the antennal joints $17 + 53 + ? + ?$, 1st joint $0.81 \times$ as long as diatone, 2nd joint $1.7 \times$ as long as basal width of pronotum; hair covering of antennae smooth, 1st joint with a whitish erect seta in each light knob. Rostrum extending beyond hind coxae. Hair covering of upper surface short, smooth and silvery, moreover light erect setae present. Pronotum with lateral margins distinctly insinuated in the middle; apical ring relatively broad; calli distinct and elevated, separated by a transverse depression from the basal portion of the disk; basal portion somewhat swollen in the middle, but without a distinct carina. Scutellum strongly sloping caudad in basal part, apex elevated. Claws of the *Cremnocephalus* type.
- Gülek, 1 ♀, 6. VIII. 1963. On *Juniperus phoenicea*.
- Pilophorus pusillus* RT. — nr. Bolu, many, 11. VIII. 1963. On *Amygdalus*.
- P. cinnamopterus* (KBM.) — nr. Gerede, 3 spec., 11. VIII. 1963. On *Pinus* sp. New to Turkey. Holarctic.