



Fig. 16. *Orthotylus junipericola terminalis* n. sp.: a—c right stylus from different angles; d left stylus; e sensory lobe of same of another specimen; f appendages of vesica. — *Platycranus orientalis* n. sp.: g—h left stylus; i—m right stylus from different angles. — *P. metriorhynchus* Rt. (specimen from S France): n—o left stylus; p—q right stylus from different angles. — Orig.

*Orthotylus nassatus* (F.) — nr. Bolu, 1 spec., 11. VIII. 1963. On *Amygdalus*. New to Turkey. European.

*O. diaphanus* (KBM.) — nr. Ankara, some, 10. VIII. 1963; nr. Kizilcahamam, some, 11. VIII. 1963. On *Salix* trees. New record for Turkey. Euro-Siberian.

*O. hodiernus* Lv. — Gülek, many, 6. VIII. 1963. On *Juniperus phoenicea*. Previously known from Israel.

*O. junipericola* Lv. ssp. *terminalis* n. ssp.

♂ 2.4—2.7 mm, ♀ 2.1—2.3 mm. As the nominate form, but considerably smaller; pronotum and scutellum bright green; vertex 1.6 (♂) or 2 × (♀) as broad as eye; 2nd antennal joint 0.83—1.0 × as long as basal width of pronotum; right stylus (fig. 16 a—c) distinctly curved apically, bearing numerous small teeth; left stylus with a rounded and relatively less prominent sensory lobe (fig. 16 d—e) and vesical appendages (fig. 16 f) much shorter and broader.

Gülek, 1 ♂ type, many paratypes, 6. VIII. 1963. On *Juniperus phoenicea*.

The nominate form occurs in Tunisia and another subspecies (ssp. *contractulus* Lv.) in Libya.

*O. minutus* JAK. — 50 km S of Aksaray, many, 7. VIII. 1963. Swept from *Chenopodiaceae* in a salt steppe.

*O. palustris* Rt. — Greece, Lake Bourou, many, 16. VIII. 1963. New to Greece.

*Heterotoma dalmatina* E. WGN. — Akçakoca road, 1 spec., 12. VIII. 1963.