

*Cantacader curtulus* Linnavuori*Cantacader curtulus* Linnavuori, 1977:6.

Material: Yemen: Yemen, holotype (LC).  
Range: Endemic.

**Tinginae***Naochila adenana* (Drake)*Cochlochila adenana* Drake, 1957:119.

South Yemen: Jebel Jihaf (Drake & Ruhoff 1965).  
Range: NE Africa (known from Ethiopia and the Sudan).

*Naochila yemenana* (Drake)*Cochlochila yemenana* Drake, 1957:121.

Material: Yemen: Jebel Sumara, 2.I.1938, 1 ex.; Wadi Thabad, N face of Jebel Sahir, 25–26.XII.1937, Scott & Britton, 1 ex. (BM, LC).  
Range: Endemic.

*Cysteochila phae* Drake & Ruhoff*Cysteochila phae* Drake & Ruhoff, 1961:148.

Material: South Yemen: Sheikh Othman-Lahej, 6–15.VII.1963, Linnavuori, 3 exx. (LC).  
Biology: On *Acacia*.  
Range: Eremian.

*Monosteira minutula* Montandon*Monosteira minutula* Montandon, 1897:101.

Material: South Yemen: Wadi Tiban, NW of Jebel Jihaf, 22.X.1937, Scott & Britton, 4 exx. (BM, LC).  
Range: Eremian.

*Urentius hystericellus* (Richter)*Tingis hystericellus* Richter, 1869:84.

Yemen: (Péricart 1983).  
Range: Palearctic.

**Miridae****Cylapinae***Rhinofulvius albifrons* (Reuter)*Fulvius albifrons* Reuter, 1895:137.

South Yemen: Aden (Reuter 1903).  
Range: Endemic.

**Deraeocorinae***\*Teratophylum insigne* Reuter*Teratophylum insigne* Reuter, 1884b:218.

Material: South Yemen: near Lahej, 6–15.VII.1963, Linnavuori, 1 ex. (LC).  
Biology: On *Acacia*.  
Range: Egypt, Palestine, Saudi Arabia, and the Sudanese subregion.

*\*Argyrotelaenus simoni* Reuter & Poppius*Argyrotelaenus simoni* Reuter & Poppius, 1912:16.

Material: South Yemen: Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 1 ex. (LC).  
Biology: On *Acacia*.  
Range: Eremian, known from Egypt, the Sudan, Palestine, Saudi Arabia, and Iraq.

*Deraeocoris pallens atramentarius* Linnavuori*Deraeocoris pallens atramentarius* Linnavuori, 1975a:6.

Material: Yemen: Ju Amlah ca 26 km NW of Sa'dah, 1950 m, IX.1950, Borri, Lanza & Poggesi, 1 ex. (MF); Ertil near San'a, 24.II.1938, Scott & Britton, holotype and many paratypes; San'a, 10–15.X.1937, Rathjens, many paratypes (BM, LC).  
Range: Eremian, also known from Saudi Arabia and Iraq.

*Deraeocoris dentifer* Linnavuori*Deraeocoris dentifer* Linnavuori, 1975a:6.

Material: Yemen: Beit Baus near San'a, 28.I.1938, Scott & Britton, holotype (BM).  
Range: Endemic.

*\*Deraeocoris ostentans* Stål*Deraeocoris ostentans* Stål, 1855:37.

Material: Yemen: several exx. from San'a, 2–9.X.1937, Rathjens; Usaifira near Ta'izz, 13.XII.1937; Wadi Dhahr near San'a, 21.I.1938, Scott & Britton (BM, LC).  
Range: Widely distributed in tropical Africa.

*\*Deraeocoris alluaudi* Poppius*Deraeocoris alluaudi* Poppius, 1919:123.

Material: South Yemen: Jebel Jihaf, X.1938, Scott & Britton, 1 ex. (BM).  
Range: East African.

*\*Stethoconus scutellaris* (Schouteden)*Apollodotus scutellaris* Schouteden, 1946:328–329.

Material: South Yemen: near Dhala, 1 ex. apparently belonging to this species (LC).  
Range: Previously known from Zaire (Rutshuru).