



Fig. 15. *Proboscidocoris pilostriatus* Odhiambo: a) penis; b) left style; c) right style.

Orthotylus indigoferae Linnavuori

Orthotylus indigoferae Linnavuori, 1975a:50–51.

Material: Yemen: Usaifira near Ta'izz, 13.XII.1937, Scott & Britton, some exx.; — South Yemen: Wadi Tiban NW of Jebel Jihaf, 22.X.1937, Scott & Britton, holotype and a paratype (BM, LC).

Biology: On *Indigofera oblongifolia*.
Range: Endemic.

**Orthotylus arabicus* Wagner

Orthotylus arabicus Wagner, 1962:215–216.

Orthotylus jordanicus Wagner, 1963:270–272.

Orthotylus halaibicus abbreviatus Linnavuori, 1975a:54 (Linnavuori 1986).

Material: Yemen: San'a, 7–8.IX.1937, Rathjens, 4 exx. (BM); — South Yemen: Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 4 exx. (LC).

Range: Eremian with an eastern distribution.

Pilophorinae

Aloea nigrifula Linnavuori

Aloea nigrifula Linnavuori, 1975a:62.

Material: Yemen: Kaulat-el-Asakeir near San'a, 9400 ft., 4.II.1938, Petrie, holotype and 6 paratypes (BM, LC).

Biology: On *Aloe*.
Range: Endemic.

Hallodapinae

Aeolocoris alboconspersus Reuter

Aeolocoris alboconspersus Reuter, 1903:17.

Carinonotus phytocoroides Lindberg, 1956:54–56.

South Yemen: Aden (Reuter 1903).
Range: Eremian.

**Hallodapus costai* (Reuter)

Laemocoris costai Reuter, 1890:257.

Allodapus longicornis Reuter, 1904:12.

Material: Yemen: Wadi Jaira, tributary of Wadi Siham, 10.III.1938, Scott & Britton, 1 ex.; — South Yemen: Al Huseini near Lahej, 27.XI.1937, Scott & Britton, 1 ex. (BM).

Range: Eremian.

**Hallodapus ochraceus* Linnavuori

Hallodapus ochraceus Linnavuori, 1961:211–212.

Material: South Yemen: Lahej-Dhala road, 13–14.VII.1963, Linnavuori, 3 exx. (LC).

Biology: At lamp in sandy habitats.

Range: Eremian, previously known from Saudi Arabia and the Sudan.

Systellonotus usaifirae Linnavuori

Systellonotus usaifirae Linnavuori, 1972b:42.

Material: Yemen: Usaifira near Ta'izz, 13.XII.1937, Scott & Britton, holotype (BM).

Range: Endemic.

Laemocoris trimaculatus Linnavuori

Laemocoris trimaculatus Linnavuori, 1964:326.

Material: South Yemen: Lahej-Dhala road, holotype (LC).

Range: Endemic.

Phylinae

Stirophyllus aristidae Wagner

Stirophyllus aristidae Wagner, 1965:232–234.

Material: Yemen: San'a, 7900 ft., 2–9.X.1937, Rathjens, 1 ex. (BM).

Range: Eremian.

Stirophyllus lineatus Linnavuori

Stirophyllus lineatus Linnavuori, 1971b:132.

Material: South Yemen: Lahej-Dhala road, 13–14.VII.1963, Linnavuori, holotype and several paratypes (LC).

Biology: In dunes on *Aeluropus litoralis*.
Range: Endemic.