



Fig. 1. *Alloemimus kurdus* Hoberlandt. a: female head and thorax in lateral view; b–c: right style; d–e: left style; f–g: theca in lateral and in dorsal view; h: vesica. — *A. unifasciatus* (Reuter) (ex from Tunisia, Oued Zeroud). i: left style; j: theca in lateral view; k: apex of vesica.

Tribe Phylini

Nasocoris albipennis Lindberg, 1939
Fig. 2a–d

Material: Many exx from Abu Ghair, 15–16.IV.1980; Tharthar lake, 23.VIII.1979; Waqisah, 1.V.1980, Linnavuori.

Biology: On *Ephedra alata* in semideserts.

Distribution: Eremian, known from Egypt, Saudi Arabia and Israel.

Nasocoris serratus Linnavuori, 1984

Material: As Salman-Takhadid, ♂ holotype, several paratypes, 17–18.IV.1980, Linnavuori, in coll. Linnavuori (AMNH).

Biology: On *Ephedra foliata* in a little gravelly wadi.

Distribution: Endemic.

Macrotylus (Macrotylus) perdictus Kiritschenko, 1938

Material: Aqra, 8 exx, 12.V.1981; Kalak, 1 ex, 13.V.1981, Linnavuori.

Biology: On *Echinops* in hilly steppes of northern Iraq.

Distribution: Syrio-Anatolian, recorded from Cyprus, Israel, Syria, Turkey, Iran, and southern Russia.

Macrotylus (Alloeonycha) ancyranus
Seidenstücker, 1969

Material: Gully Ali Beg, 2 exx, 13.VI.1980, Linnavuori.

Biology: In a hilly habitat. In Turkey found on *Salvia*.

Distribution: Anatolian, previously known from Turkey.