



Fig. 2. *Nasocoris albipennis* Lindberg. a: right style; b: left style; c: theca; d: vesica. — *Oncotylus basicornis* Horvath. e: female head and pronotum; f: 1st antennal joint; g: male 3rd antennal joint; h: claw; i: right style; j-k: left style from side and from above; l: theca.

Macrotylus (Alloeonycha) dahukanus
Linnavuori, 1988

Material: Brozah, holotype male, a female paratype, 13.V.1981; Darbandikhan, 1 paratype, 13.V.1980, Linnavuori, in coll. Linnavuori (AMNH).

Biology: In mountain meadows.

Distribution: Endemic.

Oncotylus (Cylindromelus) setulosus
(Herrich-Schäffer, 1837)

Material: Qora, 9 exx, 13.VI.1980, Linnavuori.

Biology: In a mountain meadow. On *Centaurea* (Wagner 1975a:249).

Distribution: Holomediterranean.

Oncotylus (Oncotylus) basicornis Horvath, 1901
Figs. 2e-l, 3a

Material: Shaykh Addy, 5 exx, 11.V.1981, Linnavuori.

Description in Wagner 1975a:253-254 is completed with the following data: Body 4.0 (♂) or 3.2 (♀) × as long as basal width of pronotum. Ocular index 2.1 (♂), 2.16-2.36 (♀). Proportions between antennal segments 11:47:33:17 (♂), 12:43:28:17 (♀), 2nd joint 2.1 (♂) or 1.72-1.83 (♀) × as long as diatone, 1.31 (♂) or 1.02-1.16 (♀) × as long as basal width of pronotum; under surface of 3rd antennal segment in ♂ with long hairs. Hind tibia about 2.45 (♂) or 1.8-2.2 (♀) × as long as basal width of pronotum.

Male genitalia in Figs. 2i-1, 3a.

Biology: In mountain meadows.

Distribution: Syrio-Anatolian; recorded from Syria (Akbes) and Turkey (Gaziantep).