

Pronotum green, finely rugulose, width at base equal to 2.8 times length; surface covered with erect black setae, intermixed with pale, recumbent, silky hairs; calli small, indistinct; a single long seta present at base of each anterior angle; posterior angles acute, rounded. Scutellum green to bright yellow, slightly raised; mesoscutum broadly exposed; surface finely rugulose, set with erect fuscous hairs.

Hemelytra green, subtranslucent, shining; set with erect fuscous setae interspersed with pale, recumbent, silky hairs; fuscous hairs more dense on cuneus; membrane uniformly fumate, veins dull green.

Venter dull green, abdomen darker; rostrum pale green, tip black, length 1.32 mm, reaching beyond hind coxae. Legs green to yellowish, covered with very short, erect, fuscous hairs; femora with scattered longer pale hairs; tibial spines stout, fuscous; apices of tibiae and terminal tarsal segments infuscated. Abdomen with fine golden hairs; genital segment lacking a tergal process; left clasper slender, curved, pointed at tip; right clasper weakly sinuate, blunt (see Fig. 1).

Female: Similar to male in structure, pubescence, and coloration; length 3.48 mm; maximum width 1.40 mm.

Material examined.—Holotype, male, and allotype: ARIZ., Gila Co., Rye, 1120 m (3500'), IV-18-82, D. A. & J. T. Polhemus (JTP). Paratypes: 13 ♂, 23 ♀, taken with types on *Juniperus monosperma* (One Seed Juniper); 3 ♂, 5 ♀, ARIZ., Cochise Co., Portal, 1526 m (4770'), IV-28-81, D. A. & J. T. Polhemus (JTP), on *Juniperus monosperma*.

Etymology.—The name *geronimo* is a noun in apposition, and refers to the Apache chief who for years eluded the U.S. cavalry amidst the Arizona mountains.

Discussion.—*Dichaetocoris geronimo*, n. sp. has a general appearance similar to that of *D. coloradensis* Knight, but may be separated from this and all other *Dichaetocoris* species by its blunt, relatively unmodified right clasper (see Fig. 1). The left clasper is thin and curving, ending in a point, and very similar in shape to those of *D. pinicola* Knight and *D. anasazi* Polhemus. The types were taken on juniper in the company of several species of *Dichroscytus*, a mirine genus with which these bugs may be easily confused.

Dichaetocoris mojave, NEW SPECIES

Description.—Male: Small, ovate; length 3.68 mm; maximum width 1.40 mm. Coloration uniformly golden brown.

Head golden brown, eyes black; head oriented vertically, broad; frons with erect fuscous setae intermixed with pale, recumbent, silky hairs; gula with patch of fine, pale, erect hairs; vertex set with erect fuscous hairs, width subequal to twice the dorsal width of an eye. Antennae pale brown to golden; segments I–IV with short, black, bristle-like hairs; segment I bearing three long, stout, fuscous hairs; segments III and IV with scattered erect pale hairs; lengths of antennal segments I–IV (in mm): 0.24; 1.08; 0.68; 0.32.

Pronotum golden brown, trapezoidal, width at base equal to twice its length; surface finely rugulose, set with erect fuscous setae intermixed with pallid silky hairs; calli small, indistinct; a single long seta present near each anterior angle; posterior angles rounded. Scutellum golden brown, flat, mesoscutum broadly exposed; surface set with erect fuscous hairs.

Hemelytra golden brown, shining, subtranslucent; set with erect fuscous setae