



Fig. 3. *Acraethus punctiger* (Linnavuori): a) male head and pronotum in lateral view; b–d) right style; e) apex of right style from above; f–h) left style in different views; i) apex of left style from above (dry mount). — *A. ocellaris* sp. n.: j) male head and pronotum, lateral view; k) hind leg; l) claw; m–o) right style; p) apex of right style from above; q–r) left style; s) left style from above (dry mount); t) aedeagus, lateral view.

mesocorium nearly impunctate, cuneus finely punctate. Propleura densely, meso- and meta-pleura sparsely and finely punctate. Legs long and gracile. Proportions between hind tarsomeres 7:12:12.

Male genitalia in Figs. 2v, 3b–i. Middle tooth on apical margin of right style small. Apical process of hypophysis of left style relatively gracile.

Distribution: West Sudanese.

Acraethus ocellaris sp. n.

Figs. 3–4

Material: Nigeria: B Pl St., Wamba-Lafia, ♂ holotype, 30.VIII.1973, Linnavuori. Numerous paratypes from Ivory Coast: Gouméré, 19.IX.1973; Foro-Foro, 23–28.IX.1973; Touba, 23.X.1973, Linnavuori. Togo: Anié, 11–12.IX.1973;

Atakpamé-Bena, 12.IX.1973; Langabou, 12.IX.1973; Nuatja, 11.IX.1973, Linnavuori. Benin: Tschaourou, 6.IX.1973, Linnavuori. Nigeria: N C St., Kagoro forest, 29–30.VIII.1973; NE St., Serti, 20.VIII.1973; B Pl St., Gangare Kibo, 28.VIII.1973, Gudi, 7.VIII.1973, Wamba, 7.VIII.1973, Wamba-Lafia, 30.VIII.1973; SE St., Ikom-Obudu, 23.VI.1973, Obudu cattle ranch, 16–18.VI.1973; E C St., Norcap near Abakaliki, 29.VI.1973, Linnavuori. Types in coll. Linnavuori.

Length 2–3 mm. Like the preceding species, but head black, 1st antennal joint blackish with pale base, 2nd uniformly pale yellowish, joints 3 and 4 slightly embrowned. Membrane of elytra with large roundish hyaline spot on base of lateral margin, median margin with another hyaline area. Femora black.

Head about 0.71 × as broad as basal width of pronotum, in apical view as broad as high. Eyes small, in ♂ only slightly larger than in ♀, ocular