



Fig. 5. *Acraetheus albipes* (Motschulsky) (ex from Calcutta): a–d) right style; e) apex of right style from above; f–g) left style. — *Nanniella pallidiceps* Linnavuori: h) male head and pronotum, dorsal view. — *N. gracilis* Linnavuori: i) male head and pronotum, dorsal view; q) right style; r) left style. — *N. chalybea* Reuter: j) male head and pronotum, dorsal view; k) female head, apical view; l) apex of prothorax in lateral view (co = collar, c = coxa); s) claw; t–v) left style. — *N. reuteri* Poppius: o) elytron (cu = cuneus); p) female head and pronotum, dorsal view. — *N. palustris* Linnavuori: m–n) female and male head in lateral view.

basal width of pronotum, in apical view $1.14 \times$ as broad as high; eyes small, ocular index 2.0. Rostrum extending to hind coxae. Pronotum remarkably elongate, $1.43 \times$ as broad as long (total length), lateral margins distinctly insinuated; collar broadish, disk convex, coarsely punctate. Puncturing on scutellum and elytra also dense and coarse, mesocorium with impunctate apical patch, cuneus apically impunctate.

Male genitalia in Fig. 8d–e.

Distribution: Zaire (Katanga).

***Nanniella gracilis* Linnavuori**

Fig. 5

Nanniella gracilis Linnavuori 1975:49.

Types: The Sudan, Equatoria, near Gilo, ♂ holotype and some paratypes, 18–24.III.1963, Linnavuori, in coll. Linnavuori.

Material: Cameroon: Dimako, 5 exx, 12–13.VI.1973, Linnavuori.

Length 3.5–3.75 mm. Shiny black. Eyes brownish. Antennae with 1st and 2nd joints and base of 3rd narrowly yellowish, other parts black. Cuneus apically broadly pale; membrane dark with paler spots. Femora yellow-brown, on anterior surface somewhat infumed. Tibiae black, sometimes ornamented with pale subapical rings. Tarsi bicolored, 1st and 2nd joints yellowish, 3rd joints contrastingly black.

Body large and gracile, about $3.8 \times$ as long as broad at base of pronotum. Hair covering grayish. Ocular index 2.1–2.2. Antennae long and gracile, nearly as long as body; proportions between joints 10:34:22:20 (♂), 23:70:58:58 (♀), 2nd joint 1.5 (♂) or 1.2 (♀) \times as long as basal width of pronotum. Rostrum extending to hind coxae. Pronotum narrow, $1.4 \times$ as broad as long (total