



Fig. 42. Dorsal view of *Felisacodes bryocorina* (Poppius): a), *Zanchius alatanus* Hoberlandt: b), *Z. breviceps* (Wagner): c). — After Carvalho, Dutra & Becker 1960, van Doesburg 1984 and Hoberlandt 1955.

clavus in inner basal angle of corium blackish brown, apex of cuneus infumed; membrane brownish smoky, apex and the large cell whitish hyaline. Under surface of head and thorax and lateral margins and apex of venter blackish; anterior part of pronotum and middle of venter yellowish. Legs pale yellowish or greenish.

Body very gracile, nearly 4 × as long as broad. Hair covering long and pale. Head 0.65–0.72 × as broad as basal width of pronotum; ocular index ♂ 1.11–1.20, ♀ 1.50–1.55. Antennae very long and gracile, finely haired; proportions between joints 47:121:7 :52, 1st joint tapering apicad, 1.07–1.12 × as long as diatone, 2nd 1.95 (♂) or 1.73 (♀) × as long as basal width of pronotum. Rostrum long, extending to base of venter. Pronotum 1.5–1.6 × as broad as long in middle, apical margin rugose, callal area globose and glabrous; posterior part of disk convex, transversely wrinkled and rugose. Elytra narrow, nearly 7 × long as broad (width measured at tip of clavus). Propleura rugose. Legs very long and gracile, hind tibia about 2.6 × as long as basal width of pronotum.

Male genitalia in Fig. 41j–s.

Biology: On undergrowth in mountain forests. Host: *Plectranthus fruticosus* (Schuh 1974:41).

Distribution: Widely distributed in tropical Africa. Originally described from Chirinda in S Rhodesia.

Genus *Zanchius* Distant

Zanchius Distant 1904:477. Type species: *Z. annulatus* Distant.

Uzeliella Poppius 1911:31. Type species: *U. flavovirens* Poppius (Carvalho 1952:79).

Poppiella Bergroth 1911:188, new name for *Uzeliella* Poppius 1911 (nec *Uzeliella* Baguel 1908, Thysanoptera). Type species: *Uzeliella* Poppius (Carvalho 1952:79).

Zonodorus Distant 1909b:522. Type species: *Z. gubernator* Distant (Carvalho 1958:133).

Habrocoris Wagner 1951:153–155. Type species: *H. breviceps* Wagner (Linnavuori 1964:329).

Diagnosis: Gracile green species. Head short and quadrate; eyes separate from anterolateral corners of pronotum. Pronotum small, lateral and basal margins insinuated, mesoscutum broadly visible. Pygofer conical. Right style small.