

Zheng and Wang, 1983 [Entomotaxonomia 5: 48, 57], from China, a junior primary homonym of *Lygus pratensis pubescens* Reuter, 1912 [Ofv. Finsk. Vet. Soc. Forh. 54A(7): 37], = *Lygus rugulipennis* Poppius, 1911, from the Palearctic.

Lygocoris wangi nom. n. is proposed for *Lygocoris (Apolygus) signatus* Zheng and Wang, 1983, *Lygus (Apolygus) signatus* Zheng and Wang, 1983 [Entomotaxonomia 5: 48, 56], from China, a junior primary homonym of *Lygus* [now *Pinalitus*] *cervinus* form *signata* Stichel, 1930 [Illstr. Bestimmungstabellen Deut. Wanzen 6, 7: 181], from Europe.

Lygocoris zhengi nom. n. is proposed for *Lygocoris (Apolygus) marginatus* Zheng and Wang, 1983, *Lygus (Apolygus) marginatus* Zheng and Wang, 1983 [Entomotaxonomia 5: 48, 56], from China, a junior secondary homonym of *Phytocoris marginatus* Zetterstedt, 1838 [Insecta Lapponica descripta 1: 272], = *Lygocoris rugicollis* (Fallen, 1807), from the Palearctic.

Orthotylus costai nom. n. is proposed for *Orthotylus clarensis* (Carvalho, 1990), *Melanotrichus clarensis* Carvalho, 1990 [An. Acad. Brail. Cienc. 62: 395], a junior secondary homonym of *Orthotylus clarensis* Carvalho and Costa, 1990 [An. Acad. Brasi. Cienc. 62: 310], both from Brazil [see discussion of *Orthotylus* under New Combinations for this and other species of *Orthotylus*].

Orthotylus harryi nom. n. is proposed for *Orthotylus pallens* (Knight, 1968), *Melanotrichus pallens* Knight, 1968 [Brigham Young Univ. Sci. Bull., Biol. Ser., 11(3): 125], from western USA, a junior secondary homonym of *Orthotylus pallens* (Matsumura, 1911), *Calocoris pallens* Matsumura, 1911 [J. Coll. Agric. Tokyo Imp. Univ., Sapporo 4: 39], from Sakhalin Island.

Orthotylus hazeltoni nom. n. is proposed for *Orthotylus uniformis* (Knight, 1968), *Melanotrichus uniformis* Knight, 1968 [Brigham Young Univ. Sci. Bull., Biol. Ser., 11(3): 128], from western USA, a junior secondary homonym of *Orthotylus uniformis* Van Duzee, 1916 [Proc. California Acad. Sci., 4th ser., 6(5): 100], from California.

Orthotylus leonardi nom. n. is proposed for *Orthotylus elongatus* (Kelton, 1980) *Melanotrichus elongatus*, Kelton [Can. Entomol. 112: 338], from Canada, a junior secondary

homonym of *Orthotylus elongatus* Wagner, 1965 [Reichenbachia 6: 67], from Cyprus.

Orthotylus subtropicalis nom. n. is proposed for *Orthotylus missionensis* (Carvalho and Carpintero, 1986), *Melanotrichus missionensis* Carvalho and Carpintero, 1986 [Rev. Brasileira Biol. 46: 619], a junior secondary homonym of *Orthotylus missionensis* Carvalho 1985 [Rev. Brasileira Biol. 45: 280], both from Argentina.

Orthotylus josei nom. n. is proposed for *Orthotylus cuneatus* Carvalho, 1985 [Rev. Brasileira Biol. 45: 277], from Argentina, junior primary homonym of *Orthotylus cuneatus* Van Duzee, 1916 [Proc. California Acad. Sci. (4)6: 117], from California.

Phoenicocoris knighti nom. n. is proposed for *Phoenicocoris pini* Knight, 1968, *Lepidopsallus pini* Knight, 1968 [Brigham Young Univ. Sci. Bull., Biol. Ser., 9: 53; see Stone-dahl, 1990, Bull. Amer. Mus. Nat. Hist. 198: 14], from western USA, a junior secondary homonym of *Atractotomus pini* Fieber, 1864 [Wien. Entomol. Monatschr. 8: 224], = *Phoenicocoris obscurellus* (Fallen, 1829), from the Palearctic.

Phytocoris femoratus nom. n. is proposed for *P. femoralis* Fieber, 1861 [Eur. Hem.: 260] from the Mediterranean, a junior primary homonym of *P. femoralis* Lucas, 1849 [Expl. Alg.: 82] = *Calocoris fulvomaculatus* (De Geer, 1773), Palearctic.

Phytocoris jorigtooi nom. n. is proposed for *Phytocoris desertorum* Nonnaizab and Jorigtoo, 1992 [Zool. Res. 13: 318, 322], from China, a primary homonym of *Phytocoris desertorum* Reuter, 1900 [Ofv. Finsk. Vet. Soc. Forh. B 42: 251], from North Africa.

Phytocoris leucopterus nom. n. is proposed for *P. albipennis* Reuter, 1904 [Ofv. Finsk. Vet. Soc. Forh., 47(4): 3] from Israel, a junior primary homonym of *P. albipennis* Fallén, 1829 [Hem. Suec., p. 107] = *Plagiognathus albipennis* (Fallén, 1829), Holarctic.

Phytocoris marmoratus Blanchard, 1852, in Gay [Hist. Fis. Polit. Chile, 7: 194] is the valid name for *P. chilensis* Carvalho, 1955 [Rev. Chilena Entomol. 4: 225] from Chile. **Note:** In renaming the preoccupied *P. adspersus* Spinola, 1852, Carvalho (1955) gave it a new name *P. chilensis*, instead of using the synonymic name *P. marmoratus*.

Phytocoris nonnaizabi nom. n. is proposed for *Phytocoris procerus* Nonnaizab and Jo-