

Usinger (CAS); 1 ♂, Graham Co., Stockton Pass, Pinaleno Mts., elev. 5,200–5,500 ft, June 1–2, 1983, R. T. Schuh and G. M. Stonedahl colls., collecting at night on white sheet attracted by mercury vapor trap (AMNH); 14 ♂♂, 12 ♀♀, Mohave Co., Hualapi Mts., SE Kingman, T20W R15W, 4,000–6,400 ft, June 9–10, 1983, R. T. Schuh, M. D. Schwartz, and G. M. Stonedahl, taken on *Quercus gambelii* Nutt. (Fagaceae) (AMNH; 4 USNM). **Colorado:** 5 ♀♀, Douglas Co., Perry Park, VII-8-82, J. T. Polhemus, on *Quercus gambelii* (Gambel oak) (JTPC); 3 ♂♂, 4 ♀♀, Garfield Co., 10 mi E. Glenwood Springs, VI-22-82, J. T. Polhemus, on *Quercus gambelii* (Gambel oak) (JTPC); 2 ♀♀, Garfield Co., W Evacuation Creek, 11 mi SE CO St. Border, on UT Rt. 45, el. 6,800 ft, July 9, 1981, M. D. Schwartz, taken on *Quercus gambelii* Nutt. (Fagaceae) (AMNH); 3 ♂♂, Grand Co., nr. Taylor Flats, VII-4-80, J. T. and D. A. Polhemus (JTPC); 5 ♀♀, Jefferson Co., Deer Creek Cyn., VII-27-82, J. T. Polhemus, on *Quercus gambelii* (Gambel oak) (JTPC); 1 ♀, Montrose Co., 18 mi E. of Naturita on Colo. 90, VII-7-80, J. T. and D. A. Polhemus (JTPC). **New Mexico:** 8 ♂♂, same data as for holotype (JTPC, NMNH). **Utah:** 3 ♂♂, 15 ♀♀, Carbon Co., entr. Price Cyn Recr. Area, 8 mi NW Helper, rt. 50/6, T12E R9E, 8,000 ft, July 9, 1982, M. D. Schwartz, on *Quercus gambelii* Nutt (AMNH); 32 ♂♂, 10 ♀♀, Garfield Co., Mt. Hillers at Starr Springs Cmpgrnd, T34S, R11E, 6,300 ft, June 18, 1983, R. T. Schuh and M. D. Schwartz colls., at mercury vapor light (AMNH; 6 USNM); 1 ♀, Salt Lake Co., mouth of Little Cottonwood Can., on UT St. Rt. 210, elev. 5,800 ft, June 28, 1981, M. D. Schwartz, on *Quercus gambelii* Nutt. (AMNH); 1 ♀, Sanpete Co., Cottonwood Creek, 4.7 mi NE Fairview on 31, 0.5 mi. W Nat'l For. Rd, 7,000 ft, July 12, 1981, M. D. Schwartz, on *Quercus gambelii* Nutt. (AMNH); 2 ♀♀, Sevier Co., Clear Creek Narrows Summit on Rt. 4, 2,244 m, July 15, 1980, R. T. Schuh and G. M. Stonedahl, on *Quercus* sp. (AMNH); 4 ♀♀, Sevier Co., 1.3 mi E Hwy 25 on Mytoge Mtn. Rt., VII-16-1980, 8,600 ft, M. D. Schwartz, labeled as on *Artemisia* sp. (AMNH); 3 ♂♂ (one nymph), Utah Co., Provo Canyon, rt. 92, 1 mi W Sundance Ski Resort, el. 8,000 ft, T4S R3E, July 4, 1982, M. D. Schwartz, on *Quercus gambelii* Nutt (Fagaceae); 1 ♀, Wasatch Co. Dock Flat, 1 mi NE of Ut St. rt. 40, T2S W12W, Sec. 9, elev. 8,000 ft, Aug. 14, 1982, M. D. Schwartz, on *Quercus gambelii* Nutt (AMNH); 7 ♀♀, Wasatch Co., 5 mi W Duchesne Co. line on Ut St. rt. 40, Deep Cyn. T3S R9W, 7,000 ft, July 5, 1981, M. D. Schwartz, on *Quercus gambelii* Nutt (AMNH); 5 ♀♀, Vivian Park, VII-7-22, E. P. Van Duzee (CAS).

Genus *Icodema* Reuter, 1875

Icodema Reuter 1875: 97. Type species: *Plagiognathus infuscatus* Fieber 1861. Monotypic.

Diagnosis. This genus is recognized in the Palearctic by the keeled male genital capsule, the lack of dark spots at the bases of the tibial spines, and the pallid to pale greenish white overall coloration, with only the first antennal segment infuscated and, according to Wagner (1975), with the membrane dark spotted. Autapomorphic for the genus are the very slender S-shaped vesica, with the secondary gonopore well back from the apex, and laterally notched phallosome.

Description. Length 3.70 to 4.70 mm. Elongate, oval. Overall coloration pale yellow. Head broader than long, slightly wider at base than anterior margin of pronotum;