

Currently *Heterolygus* id.

**elegans* Reuter, 1904 (*Stenodema*). LT (GC48): ♀, pn, Museum Paris, Mou-Pin, A. David, 1870/ Tibet (Mou-Pin), David 1689-70/ Type [rl]/ *Stenodema elegans* Reut. Typ. [hR]/ *Stenodema elegans* Reut [hR]. PLT: 1 ♂, hemelytra partly broken and glued on a separate card below, pn, first two labels as in LT/ *Stenodema elegans* Reut. ♂ [hR]; 1 ♀, pn, first two labels as in LT.

**elevatus* Fieber, 1858 (*Macrocoleus*). LT (CP52): ♀, gc, gs/ Hyères [hP, on green paper]/ Museum Paris, Coll. A. Puton, 1906. PLT (CN): 1 ♀, id. prep., gs. Currently *Macrotylus* id.

ericetorum Reuter, 1899 (*Psallus*). ST (CP55): 1 ♀, gc, Corfu [h]/ pbs/ *Psallus ericetorum* Rt. [hP].

**erythronotum* Reuter, 1902 (*Calocoris sexpunctatus* var.). LT (GC59): ♀, pn, Museum Paris, Tunisie, Carthage, J. De Gaulle, 1898/ Tunisie, Carthage, 1-4-58, de Gaulle/ v. *erythronotum* Reut. Typ. [hR]. Synonym of *C. nemoralis* (Fabricius, 1787).

**fasciaticollis* Reuter, 1903 (*Adelphocoris*). LT (GC 48): ♀, pn, Museum Paris, montagnes au nord de Pekin, A. David 80-63/ *Adelphocoris fasciaticollis* Reut. Typ. [hR]. PLT: 2 ♀, as LT, but without last label.

**femorialis* Reuter, 1899 (*Phylidea*). LT (CP53): ♂, gc, Nyons [hP]/ pbs/ *Phylidea Henschi* Reut. = *femorialis* Reut. sec. Reut. [hR]. Synonym of *Psallus henschi* (Reuter, 1888).

**fieberi* Fieber, 1858 (*Pachypterna*). LT (CN): ♀, pn, gs. PLT: 1 ♀, gc [was pinned initially], without labels.

**fieberi* Fieber, 1864 (*Psallus*). LT (CN): ♂, gc [card of subquadrate form characteristic of Douglas and Scott's specimens], gs. Currently *Psallodema* id.

**fieberi* Fieber, 1858 (*Tragiscus*). LT (CN): ♂, gc, gs. PLT: 1 ♂, as LT. Currently *Tragiscocoris* id.

flavipes Reuter, 1899 (*Sthenarus*). ST (CP54): 1 ♀, Mékalis (Oran), L. Bleuse, 16-5.1896/ pbs/ *Sthenarus flavipes* Rt. [hP]. Currently *Psallus* id. - Note. Wagner (1960a: 9) referred to a specimen from ZMHU as "Type". Although his wording does not constitute designation of a LT, it is preferable to designate as LT the specimen examined by him.

**forelii* Fieber, 1858 (*Orthops*). LT (CP45): ♀, gc, gs/ Type [rl].

**freyi* Puton, 1887 (*Capsus*). LT (CP46): ♀, pn, ♀/ Type [rl]/ *Capsus Freyi* Put. [hP]. PLT: 1 ♀, pn, Sicile. Synonym of *Deraeocoris flavilinea* (Costa, 1862).

**fulvipes* Reuter, 1901 (*Opisthotaenia*). LT (CP53): ♂, gc, Amasia [hP]/ pbs/ *Opisthotaenia fulvipes* Rt. [hP].

fulvus Fieber, 1836 (*Miris*). ST? (CP41): 1 ♀, pn, gs. Synonym of *Stenodema virens* (Linnaeus, 1767).

fulvus Reuter, 1875 (*Plagiognathus*). ST? (CP54): 1 ♂, gc, Tours [hP]/ Reuter [hP]. Synonym of *Nigrocapillocoris ochraceus* (Scott, 1872).

**funestus* Reuter, 1903 (*Adelphocoris*). LT (GC48): ♂, pn, Museum Paris, Mou-Pin, A. David 1870/ Tibet (Mou-Pin), David 1689-70/ *Adelphocoris funestus* Reut. Typ. [hR]. PLT: 2 ♀, as LT but without last label.

**fuscescens* Stichel, 1933 (*Orthotylus rubidus* f.). LT (CP50): ♀, gt, La Nouvelle/ pbs/ *rubidus* var. ♂ Reut. [hR]. Synonym of *O. salsolae* Reuter, 1875 (new synonymy). - Note. The name f. *fuscescens* Stichel, 1933 (later replaced because of homonymy by f. *bicolor* Stichel, 1957) was given to *O. rubidus* var. ♂ of Reuter (1883: 372). Reuter stated that var. ♂ is similar in

pubescence to *O. salsolae* Reuter, 1875 (described from La Nouvelle and other localities and in 1883 regarded as a variety of *O. rubidus*), but differs in embrowned head and anterior part of pronotum. This characteristic agrees with the LT, in which brownish trockened muscles of head and fore half of pronotum are seen through the cuticle as a dark spot. Wagner (1942) incorrectly referred f. *fuscescens* to *O. moncreaffi* Douglas & Scott, 1867.

**fusciceps* Reuter, 1899 (*Orthopidea*). LT (CP54): ♂, gc, Djebel Amour, Sidi bou Zid, de Vauloger/ *Orthopidea fusciceps* Rt. du Reuter [h]. PLT: 2 ♀, as LT, but without last label and with pbs.

**fuscicornis* Brullé, 1839 (*Miris*). LT (GC53): ♀, mpn, Museum de Paris, Iles Canaries, Webb-Berthelot 3-41/ 3-41 [h; on underside of a blue circle; according to registration book for 1841, = specimens from Canary Islands gifted by Webb and Berthelot]/ *Macrotylus nigricornis* Fieb., det. Reuter [hR]. PLT: 3 ♀, damaged, mpn on one pin, as LT but without last label. Nomen oblitum, synonym of *Macrotylus nigricornis* Fieber, 1864 (new synonymy). - Note. Although the above specimens are not labelled by Brullé, we are convinced that they belong to the type series of *Miris fuscicornis*. Only 3 species of *Miridae* were recorded by Brullé (1839). Two are represented in the collection (box GC53): a specimen of *Calocoris nemoralis* Fabricius (recorded by Brullé as "*Miris bipunctatus* Fab.") and 4 specimens of *Macrotylus nigricornis*, which fit relatively well to Brullé's description of *Miris fuscicornis* and his note that the species is common in Canary Islands. The name *M. fuscicornis* was not used as valid and merits suppression under the plenary powers.

**fuscicornis* Reuter, 1899 (*Sthenarus*). LT (CP54): ♂, gc, Guelt-es-Stel, Mai 1897, Dr. A. Chobaut/ pbs/ *Sthenarus fuscicornis* Rt. [hP].

**fusciloris* Reuter, 1899 (*Ochrodema*). LT (CP56): ♀ (the left specimen), gc, Biskra, Mai 1894, Chobaut/ *Ochrodema fusciloris* Reut. n. g. et sp. [hR]/ Lectotype [rl]. PLT: 1 ♀ (the right specimen on the same card). Synonym of *Maurodactylus nigrigenis* (Reuter, 1890). - Note. Published designation of LT unknown to us.

**fusciloris* Reuter, 1878 (*Plagiognathus*). LT (CP53): ♀, gt, Vaucluse/ pbs. PLT: 1 ♀, id. prep., pbs.

**fuscinervis* Reuter, 1875 (*Atractotomus*). LT (CP 55): ♂, gt, Corse [hP]/ pbs. Synonym of *Lepidargyrus ancorifer* (Fieber, 1858).

**geniculata* Fieber, 1858 (*Brachyceroea*). LT (CN): ♂, gc, gs/ Corsica [h, on green paper]. PLT (CP49): 1 ♀, gc, gs. Currently *Dicyphus* id.

**gonoceroides* Reuter, 1903 (*Pantilius*). LT (GC48): ♂, pn, Museum Paris, Mou-Pin, A. David, 1870/ Tibet (Mou-Pin), David 1689-70/ *Pantilius gonoceroides* Reut. Typ. [hR].

**gracilis* Puton, 1874 (*Macrocoleus*). LT (CP52): ♂, gt, Balaruc [hP]/ (*Senecio viscosus*) [hP]. PLT: 1 ♀, above LT on the same pin; 1 ♂, 1 ♀, glued on one triangle, Agde [hP]. PLT?: 1 ♂, 3 ♀, gt, on 3 pins placed after LT, without labels. - Note. In CP52 id. prep. specimens are also from Cassis (not type locality), therefore origin of unlabelled specimens is problematic. Synonym of *Macrotylus atricapillus* (Scott, 1872).

guttulatus Reuter, 1894 (*Phytocoris*). LT (Wagner, 1968b: 92, as HT) (CP42): ♀, gc, Ain Draham, 1886/ pbs/ *Phytocoris guttulatus* Reut. E. Wagner [rl].

henkei Jakovlev, 1875 (*Calocoris*). ST (CP44): 1 ♀, gc