



Fig. 43. *Boopidella fasciata* Reuter: a–b) male head and thorax in dorsal and lateral view; c) head, apical view; d) hind leg; e) claw; f–g) left style; h) theca; i–j) vesica. — *Pongocoris vittatus* (Odhiambo): k–n) male and female head in dorsal and lateral view; o) hind leg. — *P. opertus* Linnavuori: p–q) male head in dorsal and lateral view; r–s) female head and thorax in dorsal and lateral view; t) right style; u–v) left style; w–x) theca; y) vestiture of hind tibia.

tions between antennal joints 8:21:13:12, 2nd joint nearly 0.9 × as long as basal width of pronotum. Pronotum nearly twice as broad as long in middle.

Male genitalia in Figs. 43t–x, 44v, 48p. Theca digitate, provided with broad trough-like lateral lamella. Vesica with long falcate apical portion.

Distribution: Only known from the southern parts of the Sudan.

***P. vittatus* (Odhiambo), comb. n.**

Figs. 36d, 43k–o, 48l–o

*Trichophthalmocapsus vittatus* Odhiambo 1959:661–663.

*Hallodapus vittatus* Schuh 1974:94.

Types studied: Uganda, Kawanda, female allotype and 1♂ paratype 19.I.1959, Odhiambo in the British Museum, 1♂ paratype, 11.XI.1959, Odhiambo, in Mus. Paris.

Length 3.2–3.4 mm. Resembling *P. opertus* but eyes with long erect setae and much larger, ocular index 0.69–0.73 (♂), 1.67 (♀), in lateral view extending to ventral margin of head even in ♀.

Male genitalia: Theca slender, provided with narrow lateral lamella. Falcate apical portion of theca short.

Distribution: Known from Uganda, Kenya and Ethiopia. 5.X.1957.

**Genus *Boopidella* Reuter**

*Boopidella* Reuter 1907:25. Type species: *B. fasciata* Reuter.

*Boopidella* Poppius 1914:45–46, Carvalho 1955:64 and 1958:162, Schuh 1974:121–122.

Opaque. Brown. Elytra with broad white transverse fascia just behind tip of scutellum.