

***L. hirsutus* sp. n.**

Figs. 46f, 52h–o

Material studied: Nigeria: N C St., Kagoro forest, male holotype, ♂ paratype, 7.VIII.1973; B Pl St., Jengre, ♂ paratype, 4.VIII.1973, Linnavuori, in coll. Linnavuori.

♂ length 4.25 mm. Opaque, Head opaquely shiny, dark reddish brown, eyes grayish. 1st antennal joint yellow-brown, base brown, 2nd joint black (other joints missing). Pronotum and scutellum black. Elytra reddish brown, with whitish ochraceous markings as indicated in Fig. 46j, pale spot near tip of clavus on corium very indistinct, cuneus dark purplish; membrane brown with a hyaline spot at tip of cuneus. Under surface reddish brown, hind coxae whitish. Femora and tibiae blackish, tarsi yellow-brown.

Body elongate. Hair covering on upper surface brownish, remarkably long, dense, erect. Head 0.64–0.66 × as broad as basal width of pronotum, rugose, with faint median depression; basal margin of vertex sharply carinate; ocular index 1.74–1.82. Proportions between antennal joints 20:88:?, 2nd joint 1.30–1.35 × as broad as basal width of pronotum. Rostrum extending to hind coxae. Pronotum moderately broadening caudad, lateral margins shallowly insinuated; collar broad, disk basally convex, strongly sloping apicad and laterad, very densely and strongly microsculptured, opaque. Scutellum shiny, apical hump bluntly conical. Elytra longer than abdomen, costal margins slightly insinuated. Legs long; femora with long dense semidecumbent hair covering; bristles on tibiae distinctly longer than cross-section of tibia.

Male genitalia in Fig. 52i–o.

***L. quinque maculatus* (Wagner)**

Fig. 40t–z

Paramimus quinque maculatus Wagner 1951:155.

Mimocapsus quinque maculatus Wagner 1953:477.

Laemocoris quinque maculatus Linnavuori 1984:38–40.

Material studied: The Sudan: Kassala, Halaib, 1♀, 20.I.1933, H. Priesner, in coll. Linnavuori.

The original description not repeated.

Distribution: The Red Sea Coast in the Sudan.

Acknowledgements. I am greatly indebted to Drs. J. André and G. Schmitz, Museum for Central Africa in Tervuren, Dr. A. Jansson, Zoological Museum of the Helsinki University, Dr. W. R. Dolling and Mrs. J. Palmer, the British Museum of Natural History, and Mrs. D. Pluot-Sigwalt, National Museum of Natural History in Paris for the opportunity to examine the type material in their respective collections. I also wish to thank Dr. B. Bolton of the British Museum for identification of the ants.

References

- Akingbohunge, A. 1975: *Miridae* (Heteroptera) of Nigeria: I. A new genus of Herdoniini with a new record and redescription of *Xenotomorpha carpenteri* Poppius. — *Bull. I. F. A. N.* 37 Sér. A (3): 637–640.
- 1979: A new genus and four new species of *Hyaliodinae* (Heteroptera: *Miridae*) from Africa with comments on the status of the subfamily. — *Rev. Zool. Afr.* 93:501–521.
- 1980: The African genera of Surinamellini (Heteroptera: *Miridae*) with the description of new species. — *Syst. Entomol.* 5:227–244.
- Bergroth, E. 1926: Über einige äthiopische *Miridae* (Hem. Het.). — *Deutsche Entomol. Zeit.* 1926:60–64.
- Carvalho, C. C. M. 1951: Five new genera and eleven new species of African *Miridae* (Hemiptera). — *Rev. Zoo. Bot. Afr.* 45(1–2): 100–115.
- 1952: On the major classification of the *Miridae* (Hemiptera). With keys to subfamilies and tribes and a catalogue of the world genera. — *An. Acad. Brasil. Ci.* 24(1): 31–110.
- 1955: Keys to the genera of *Miridae* of the world (Hemiptera). — *Bul. Mus. Goeldi* 11(2): 1–151.
- 1958: *Catálogo dos Mirideos do Mundo II.* — *Arq. Mus. Nac. Rio de Janeiro* 45:1–216.
- China, W. E. 1925: Additional notes on the Heteroptera of Rodrigues. — *Ann. Mag. Nat. Hist.* (9)15:163–165.
- 1932: A new species of *Trichophthalmocapsus* Poppius (Hemiptera, Heteroptera, Capsidae) from Kenya. — *Ann. Mag. Nat. Hist.* 10(10): 594–597.
- 1944: New and little known West African *Miridae* (Capsidae) (Hemiptera-Heteroptera). — *Bul. Entomol. Res.* 35(2): 171–191.
- China, W. E. & Carvalho, J. C. M. 1951: A new genus of *Hallodapini* and two new species of the genus *Nicostratus* (*Miridae*, *Phylinae*). — *Ann. Mag. Nat. Hist.* (12)4:1121–1125.
- DeLattre, R. 1950: Description de nouveaux *Miridae* africains (Hem.). — *Bull. Soc. Entomol. France* 55(10): 151–153.
- Distant, W. L. 1904: The fauna of British India, including Ceylon and Burma. *Rhynchota*, vol. 2 (Heteroptera): 421–488. — Taylor & Francis, London.
- 1909: Descriptions of Oriental *Capsidae*. — *Ann. Mag. Nat. Hist.* (8)4:440–454.