

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

Painite ($\text{CaZrBAl}_9\text{O}_{18}$): A second source in Myanmar and metasomatic origins

G.R. ROSSMAN¹, S. NAUNG², G.E. HARLOW³, AND J. HUNT^{1,3,4}

¹ Div. Geological & Planetary Sciences, California Institute Tech., Pasadena, CA 91125-2500 (grr@gps.caltech.edu)

² Mogok, Myanmar (drsaaw@myanmar.com.mm)

³ Dept. Earth & Planetary Sciences, Amer. Museum Natural History, NY, NY 10024-5192 gharlow@amnh.org)

⁴ Earth & Space Sciences, University of California, Los Angeles (jhunt@ess.ucla.edu)

Painite, $\text{CaZrBAl}_9\text{O}_{18}$, is a rare mineral that was previously known only as crystals from the vicinity of Ongaing village in the Mogok Stone Tract, Mandalay district, northern Myanmar. Specimens of the orange-red to brownish-red painite from Ongaing number at least thirteen. It has now been found as slightly water worn crystals among alluvial spinel, corundum and zircon from Namyia (Nanyaseik), Kachin State, some 300 km to the northwest of Mogok. Minor elements observed in EMP and XRF analyses of all painites include Ti, V, Cr, Fe and Hf. Inclusions within painite include liquid CO_2 , srilankaite $[(\text{Ti},\text{Zr})\text{O}_2]$, baddeleyite (ZrO_2), a CaAl-silicate, and calcite.

The new crystals purchased in Namyia, 50 km NW of Mogaung, are pale pink and dichroic from pale orangish-pink to nearly colorless. Painite from Namyia is fluorescent under UV light. CL shows both planar and irregular growth zoning on a scale of $<10 \mu\text{m}$ banding.

Cr and V control the red to brown coloration in all samples and produce optical absorption bands near 398, 455 and 550 nm. The Namyia material analyzed to date contains nearly an order of magnitude lower concentration of these elements and is proportionally less intensely colored. Crystals from both localities share a distinctive Raman spectrum.

The Ongaing and Namyia deposits, known for rubies (corundum), are sourced from the marbles of the Mogok Belt, interpreted as late Proterozoic limestones that have experienced various metamorphic events as late as Oligocene intrusions and metasomatism. The corundum's origin is ascribed either to paleosols that were Fe^{2+} -depleted by metasomatism or to Si-depletion metasomatism followed by Ca-reaction metasomatism. Painite is consistent with these models that yield refractory oxides and calc-silicate overgrowths.