

New records of Jordanian scorpions

Zuhair Amr¹, Omar A. Abed², Thabet Al Share², Nashat Hamidan²
& Lorenzo Prendini³

1. Department of Biology, Jordan University of Science & Technology, Irbid,
Jordan.

2. The Royal Society for the Conservation of Nature, Amman, Jordan.

3. Division of Invertebrate Zoology, American Museum of Natural History,
New York, U.S.A.

ABSTRACT

Fourteen scorpion species were collected during a recent field expedition to Jordan. Family Buthidae was represented by twelve species in seven genera (*Androctonus*, *Birulatus*, *Buthacus*, *Compsobuthus*, *Hottentotta*, *Leiurus*, and *Orthochirus*) and families Diplocentridae and Scorpionidae by one genus and species each (*Nebo* and *Scorpio*) respectively. *Buthacus nigroaculeatus* and *B. yotvatensis* were recorded for the first time in Jordan.

Key words: Scorpions, Jordan, Buthidae, Diplocentridae, Scorpionidae, *Buthacus yotvatensis*.

INTRODUCTION

Although the scorpion fauna of Jordan has never been systematically surveyed, it is better known than that of many other countries in the Middle East. Thus far, 18 species and subspecies representing 10 genera in three families, Buthidae (*Androctonus*, *Birulatus*, *Buthacus*, *Buthus*, *Compsobuthus*, *Hottentotta*, *Leiurus* and *Orthochirus*), Diplocentridae (*Nebo*) and Scorpionidae (*Scorpio*) have been reported from Jordan (Vachon, 1966; Levy *et al.*, 1973; Kinzelbach 1984; Amr *et al.*, 1988; Amr & El-Oran, 1994; Stathi & Mylonas, 2001; Lourenço *et al.*, 2002, 2010; Kovařík, 2003, 2012; Amr & Abu Baker, 2004; Kovařík & Whitman, 2004; Lowe *et al.*, 2014).

During September 2013, we conducted a field trip to collect scorpions from Jarash, along the desert highway to Aqaba, Ash Shawbak, Wadi Ramm and Al Mudawwarah. The present report documents the fourteen species of scorpions collected.

Materials and Methods

Scorpions were mostly collected by ultraviolet light detection at night, although some were collected by turning stones or excavating burrows during daytime. Material examined is deposited in the Collection of Arachnida and

Myriapoda at the American Museum of Natural History, New York. Table (1) indicates localities from which materials were collected.

Table 1: Coordinates for collecting sites.

Location	N	E
Al Barah, Wadi Rum	29°32'47.7"	35°19'36.3"
Al Brerah, Wadi Ramm	29°32'08.4"	35°32'14'00"
Al Gaal (Al Ghal), Wadi Rum	29°33'54.0"	35°37'48.4"
Al Hussainiha (Hysayneiah), ca. 2 km N on Amman–Aqaba Highway	30°52'26.9"	35°59'06.7"
Al Hussainiha (Hysayneiah)–Al Qadasiah (Qadeseah/Qadessyah/Qadisiyeh) road	30°35'14.1"	35°46'29.2"
Al Khoshakhashe (Kheshkhasheh), Wadi Rum	29°29'52.1"	35°25'54.4"
Al Shaubak (Showbak/Shawbak) castle, hillside opposite	30°31'35.6"	35°33'57.2"
Al Shaubak (Showbak/Shawbak) castle, slopes below	30°31'56.9"	35°33'36.1"
Ash Shawbak Castle	30°31'35.6"	35°33'57.2"
Dana Guest House, cliffs above	30°40'27.4"	35°36'29.7"
Dibbin (Dibeen) Forest Reserve	32°14'46.9"	35°49'22.0"
Jabal Al Romman	30°41'40.0"	35°33'46.7"
King Talal Dam road	32°11'52.4"	35°50'13.4"
Makheet, Wadi Rum	29°31'27.8"	35°23'02.3"
NW of Al Mudawwarah (Modawarah)	29°19'10.6"	35°58'58.9"
NW of Al Mudawwarah (Modawarah)	29°19'15.7"	35°59'51.5"
Wadi Al Ahmar	30°41'21.6"	35°34'28.1"
Wadi Al Mohark, Wadi Ramm	29°36'15.6"	35°31'00.4"
Wadi 'Araba, Qaser Al Tlah	30°49'47.5"	35°24'38.2"
Wadi 'Araba, Wadi Al Ghwaibeh (Ghwaqibeh)	30°48'13.3"	35°24'21.6"
Wadi 'Araba, Wadi Khanzeerah (Khanzairh)	30°53'39.9"	35°25'38.2"
Wadi Daba'a, ca. 1 km W of Amman–Aqaba Highway	31°35'49.2"	35°59'27.1"
Zogreet (Zograit), Jarash–Ajloun road	32°17'03.6"	35°51'27.7"

RESULTS

Family *Buthidae* was represented by twelve species in seven genera (*Androctonus*, *Birulatus*, *Buthacus*, *Compsobuthus*, *Hottentotta*, *Leiurus*, and *Orthochirus*) and families Diplocentridae and Scorpionidae by one genus and species each (*Nebo* and *Scorpio*, respectively). *Buthacus nigroaculeatus* and *B. yotvatensis* are recorded for the first time in Jordan.

Family *Buthidae* C.L. Koch, 1837

Androctonus bicolor Ehrenberg, 1828

Fig. 1A

Material Examined: 1 ♂, 1 ♀, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Androctonus bicolor* was previously reported from only a few localities in Jordan, including Aqaba, Karak, Ma'an, Petra, Wadi Ramm (El-Hennawy,

1988; Amr & El-Oran, 1994; Kovařík & Whitman, 2004). At Ash Shawbak, it was collected from the wadi and slopes below the castle ruins.

Androctonus crassicauda (Olivier, 1807)

Fig. 1B

Material Examined: 1 ♀, Al Brerah, Wadi Ramm, 11.ix.2013, Z. Amr, N. Hamidan & T. Al Share.

Remarks: *Androctonus crassicauda* is one of the most venomous scorpion species in the Middle East. In Jordan, it is widely distributed in the eastern desert and Wadi Araba to Aqaba and Wadi Ramm (Levy & Amitai, 1980; Amr et al., 1988; Amr & El-Oran, 1994; Stathi & Mylonas, 2001, Kovařík & Whitman, 2004). It was also collected from the Mediterranean region but in low numbers. It lives in horizontal burrows or rodent burrows.

Birulatus haasi Vachon, 1974

Material Examined: 1 ♀, Al Hysayneiah, ca. 2 km N on Amman–Aqaba Highway, 7.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. --1 ♂, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Birulatus haasi* was originally described from the Tafilah area (Vachon, 1974). It was redescribed by Lourenço (1999), who mistakenly suggested that it is a cave-dwelling scorpion. This is the second collection of this species from Jordan, with an additional locality from Al Hussainiha, expanding the known distribution into more arid regions of Jordan.

Buthacus nigroaculeatus Levy et al., 1973

Fig. 1C

Material Examined: 2 juv. ♂, Wadi Al Mohark, Wadi Ramm, 11.ix.2013, L. Prendini, O. Abed & L. Al Azam. -- 3 ♂, 3 ♀, Al Barah, Wadi Ramm, 10.ix.2013, Z. Amr, N. Hamidan & L. Al Azam. -- 1 ♀, Al Barah, Wadi Ramm, 10.ix.2013, Z. Amr, N. Hamidan & L. Al Azam.

Remarks: Specimens from Wadi Ramm (Kinzelbach, 1984; Stathi & Mylonas, 2001; Kovařík & Whitman, 2004), previously referred to as *Buthacus leptochelys* (Ehrenberg, 1829), are here referred to *Buthacus nigroaculeatus*, making this the first record of the species from Jordan.

Buthacus yotvatensis Levy, Amitai & Shulov, 1973

Fig. 1D

Material Examined: 1 ♀, Wadi 'Araba, Wadi Khanzeerah, 9.ix.2013, L. Prendini, Z. Amr & L. Al Azam. -- 2 ♂, 1 ♀, Wadi 'Araba, Wadi Al Ghwaibeh, 9.ix.2013, L. Prendini, Z. Amr & L. Al Azam. -- 2 ♂, 1 ♀, Wadi 'Araba, Wadi Al Ghwaibeh, 9.ix.2013, L. Prendini, Z. Amr & L. Al Azam.

Remarks: *Buthacus yotvatensis* was collected in the sand dunes of Wadi Al Ghwaibeh and Wadi Khanzeerah, the first records of this species in Jordan. It was previously found on sand dunes of Wadi Araba on the Israeli side (Levy & Amitai, 1980).

Compsobuthus jordanensis Levy, Amitai & Shulov, 1973

Material Examined: 6 ♂, 6 ♀, 1 subad. ♀, Wadi Daba'a, ca. 1 km W of Amman–Aqaba Highway, 7.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 9 ♂, 4 ♀, Al Hysayneiah–Al Qadessyah road, 7.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Compsobuthus jordanensis* was previously reported from around Wadi Daba'a, southeast of Amman, and Al-Hasa toward Ma'an (Levy *et al.*, 1973). It appears to be widely distributed in Jordan, extending further south and to the west of the Irano-Turanian zone.

Compsobuthus levyi Kovařík, 2012

Material Examined: 1 ♂, Dana Guest House, 8.ix.2013, L. Prendini. -- 5 ♂, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 4 ♂, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Compsobuthus levyi* was previously reported from Qasr Burqu, in the eastern desert of Jordan (Kovařík, 2012). The new locality records extend its distribution further into southwestern Jordan.




Figure 1: A. *Androctonus bicolor*. B. *Androctonus crassicauda*. C. *Buthacus nigroaculeatus*. D. *Buthacus yotvatensis*. E. *Compsobuthus schmiedeknechti*. F. *Hottentotta judaicus*.

Compsobuthus schmiedeknechti Vachon, 1949

Fig. 1E

Material Examined: 8♂, 2♀, Dibeen Nature Reserve, 6.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Compsobuthus schmiedeknechti* occurs in rocky habitats in the Mediterranean region of Jordan (Levy & Amitai, 1980). Published records include Bonifica and Petra (Vachon, 1949; Kovářík & Whitman, 2004).

Hottentotta judaicus (Simon, 1872)

Fig. 1F

Material Examined: 3♀, Zogreet, Jarash-Ajloun road, 6.ix.2013, L. Prendini, Z. Amr. -- 1juv., Dibeen Forest Reserve, 6.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: *Hottentotta judaicus* was previously reported from several localities within the Mediterranean region of Jordan (Wahbeh, 1976; Kinzelbach, 1984; El-Hennawy, 1988; Amr & El-Oran, 1994). It is often associated with the *terra rossa* soil.

Leiurus jordanensis Lourenço, Modry & Amr, 2002

Fig. 2A

Material Examined: 1♂, NW of Al Mudawwarah, 10.ix.2013, L. Prendini, O. Abed & T. Al Share.

Remarks: *Leiurus jordanensis* was originally described from a desert habitat in southern Jordan composed of sandstone cliffs surrounded by flat sand dunes (Lourenço et al., 2002). It was subsequently recorded from northern Saudi Arabia (Hendrixson, 2006; Lowe et al., 2014).

Leiurus hebraeus (Birula, 1908)

Fig. 2B

Material Examined: 6♂, 10♀, 1subad.♂, 1subad.♀, 9 juv.♂, 6 juv.♀, King Talal Dam road, 6.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 2♀, 1 juv.♂, 4 juv.♀, Jabal Al Rumman, 9.ix.2013, O. Abed & T. Al Share. -- 1♂, Wadi Al Ahmar, 9.ix.2013, O. Abed & T. Al Share. -- 2♂, Dana Guest House, 9.ix.2013, L. Prendini. -- 55♂, 42♀, 8 subad.♂, 8 subad.♀, 11 juv.♂, 12 juv.♀, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 50♂, 63♀, 6subad.♂, 7subad.♀, 1 juv.♂, 3 juv.♀, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 3♂, 8♀, 2subad.♀, 1 juv.♂, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam.

Remarks: This species, previously regarded as a subspecies of *Leiurus quinquesstriatus* (Ehrenberg, 1828), is the most venomous scorpion species in Jordan (Amr et al., 1994). It is also the most common species with a wide distribution covering much of the western part of the country (Levy et al., 1970; Wahbeh, 1976; Levy & Amitai, 1980; El-Hennawy, 1988; Amr & El-Oran, 1994; Stathi & Mylonas, 2001; Lowe et al., 2014). More than a hundred specimens were collected near Ash Shawbak Castle, on the walls of caves, under bushes and in rocky areas.

Orthochirus scrobiculosus negebensis Shulov & Amitai, 1960

Fig. 2C

Material Examined: 2♂, 2♀, Wadi Daba'a, ca. 1 km W of Amman-Aqaba Highway, 7.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 1♂, 1♀, Al Hysayneiah, ca. 2 km N on Amman-Aqaba Highway, 7.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 1♀, Wadi 'Araba, Qaser, Al Tlah, 9.ix.2013, L. Prendini, Z. Amr & L. Al Azam.

Remarks: The status of this taxon is unclear. *Orthochirus* populations from Jordan, Palestine, and Sinai were initially described as *O. innesi negebensis* Shulov & Amitai, 1960 and later regarded as *O. scrobiculosus negebensis* (Levy & Amitai, 1980; Fet & Lowe, 2000). Kovářík (2004: 27) suggested these populations may be a 'separate species' while Kovářík & Whitman (2004) identified specimens from Wadi Ramm as *Orthochirus innesi* Simon, 1910. Hendrixson (2006) referred specimens from neighboring Saudi Arabia to *O. innesi*. *Orthochirus scrobiculosus negebensis* is usually found under stones and was previously reported from arid and semi-regions of Jordan, e.g., Azraq, Jerash, Madaba, Thgrat al Jubb (Wahbeh, 1976; Amr & El-Oran, 1994; Stathi & Mylonas, 2001). High population densities were previously reported in the Azraq area (Amr & El-Oran, 1994). The newly reported material was collected along the desert highway within the Irano-Turanian region and from Wadi Araba.



Figure 2: A. *Leiurus jordanensis*. B. *Leiurus hebraeus*. C. *Orthochirus scrobiculosus negebensis*. D. *Nebo hierichonticus*. E. *Scorpio kruglovi*.

Family Diplocentridae Karsch, 1880

Nebo hierichonticus (Simon, 1872)

Fig. 2D

Material Examined: 1 subad. ♀, King Talal Dam road, 6.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 1 ♂, Wadi Al Ahmar, 9.ix.2013, O. Abed & T. Al Share.

Remarks: *Nebo hierichonticus* is endemic to the Levant. In Jordan, it was previously reported from several localities including Al-Mazar al-Janubi, Al Karak Amman, Jarash, Madaba, Petra, Wadi Araba, Wadi Karak, and Wadi Ramm (Wahbeh, 1976; Levy & Amitai, 1980; Amr & El-Oran, 1994; Stathi & Mylonas, 2001; Kovářík & Whitman, 2004). This species constructs burrows and can be found under rocks and in crevices. In the Jarash area, it was found in deep crevices among large boulders and in earthen walls.

Family Scorpionidae Latreille, 1802

Scorpio kruglovi Birula, 1910

Fig. 2E

Material Examined: 1 ♂, Jabal Al Rumman, 9.ix.2013, O. Abed & T. Al Share. -- 3 ♂, Ash Shawbak Castle, 8.ix.2013, L. Prendini, Z. Amr, O. Abed, T. Al Share & L. Al Azam. -- 1 ♂, Al Barah, Wadi Ramm, 10.ix.2013, Z. Amr, N. Hamidan & L. Al Azam. -- 1 ♂, Al Khoshahashe, Wadi Ramm, 11.ix.2013, Z. Amr, N. Hamidan & T. Al Share.

Remarks: *Scorpio kruglovi* was previously reported, as *Scorpio maurus palmatus* (Ehrenberg, 1828), from Ajloun, Amman, Theban, Wadi Musa, and Wadi Ramm (Wahbeh, 1976; El-Hennawy, 1988; Amr & El-Oran, 1994; Kovářík & Whitman, 2004). The new material was collected from rocky areas around Ash Shawbak and compacted sand areas at several localities in Wadi Ramm.

References

- Amr, Z.S. & Abu Baker, M. 2004. The scorpions of Jordan. *Denisia*, 14:237–244.
- Amr, Z.S. & Al-Oran, R. 1994. Systematics and distribution of scorpions (Arachnida, Scorpionida) in Jordan. *Bollettino di Zoologia*, 61:185–190.
- Amr, Z., Hyland, K., Kinzelbach, R., Amr, S. & Defosse, D. 1988. Scorpion etpiques de scorpions en Jordanie. *Bulletin de la Société de Pathologie Exotique*, 81:369–379.
- El-Hennawy, H.K. 1988. Scorpions of Jordan. *Serket*, 1:13–20.
- Fet, V. & Lowe, G. 2000. Family Buthidae C. L. Koch, 1837. In: Fet, V., Sissom, W.D., Lowe, G. & Braunwalder, M.E., Catalog of the scorpions of the world (1758–1998). The New York Entomological Society, pp. 54–286.
- Hendrixson, B.E. 2006. Buthid scorpions of Saudi Arabia, with notes on other families (Scorpiones: Buthidae, Liochelidae, Scorpionidae). *Fauna of Arabia*, 21:33–120.
- Kinzelbach, R. 1984. Die Skorpion des Naturhistorischen Museum der Stadt Mainz. Teil II: Vorderasient. *Mainzer Naturw. Archiv.*, 22:97–106.

- Kovařík, F. 2003. Eight new species of *Compsobuthus* Vachon, 1949 from Africa and Asia (Scorpiones: Buthidae). *Serket*, 8:87–112.
- Kovařík, F. 2004. Revision and taxonomic position of genera *Afghanorthochirus* Lourenço & Vachon, *Baloorthochirus* Kovařík, *Butheolus* Simon, *Nanobuthus* Pocock, *Orthochiroides* Kovařík, *Pakistanorthochirus* Lourenço, and Asian *Orthochirus* Karsch, with descriptions of twelve new species (Scorpiones, Buthidae). *Euscorpius*, 16:1–33.
- Kovařík, F. 2012. Three new species of *Compsobuthus* Vachon, 1949 from Yemen, Jordan, Israel, and Somaliland (Scorpiones: Buthidae). *Euscorpius*, 150:1–10.
- Kovařík, F. & Whitman, S. 2004. Cataloghidel Museo di Storia Naturaledell' Università di Firenze-Sezione di Zoologia «La Specola» XXII. Arachnida Scorpiones. Tipi. addenda (1988–2004) e checklist della collezione (Euscorpiinaeesclusi). *Attidella Società Toscana de Scienze Naturali –Memorie serie B*, 111:103–119.
- Levy, G., Amitai, P. & Shulov, A. 1970. *Leiurus quinquestriatus hebraeus* (Birula, 1908) (Scorpiones: Buthidae) and its systematic position. *Israel Journal of Zoology*, 19:231–242.
- Levy, G., Amitai, P. & Shulov, A. 1973. New scorpions from Israel, Jordan and Arabia. *Zoological Journal of the Linnean Society*, 52:113–140.
- Levy, G. & Amitai, P. 1980. Fauna Palaestina. Arachnida I: Scorpiones. The Israel Academy of Science and Humanities. Jerusalem.
- Lowe, G., Yağmur, E.A. & Kovařík, F. 2014. A review of the genus *Leiurus* Ehrenberg, 1828 (Scorpiones: Buthidae) with description of four new species from the Arabian Peninsula. *Euscorpius*, 191:1–129.
- Lourenço, W.R. 1999. On the phylogenetic position of the genus *Birulatus* Vachon, 1973 (Scorpiones, Buthidae) and redescription of *Birulatus haasi*. *Zoology in the Middle East*, 18:109–113.
- Lourenço, W.R. 2002. Further morphological considerations on the genus *Birulatus* Vachon (Scorpiones, Buthidae), with the description of a new species from Israel. *Revista Iberica de Aracnologia*, 6:141–145.
- Lourenço, W.R., Modrý, D. & Amr, Z. 2002. Description of a new species of *Leiurus* Ehrenberg, 1828 (Scorpiones, Buthidae) from the South of Jordan. *Revue Suisse de Zoologie*, 109:635–642.
- Lourenço, W.R., Yağmur, E.A. & Duhem, B. 2010. A new species of *Buthus* Leach, 1815 from Jordan. *Zoology in the Middle East*, 49:95–99.
- Stathi, I. & Lourenço, W. 2003. Description of a new scorpion species of the genus *Birulatus* Vachon, 1974 (Scorpiones, Buthidae) from Syria. *Zoology in the Middle East*, 30:105–110.

- Stathi, I. & Mylonas, M. 2001. New records of scorpions from the central-eastern Mediterranean area: biogeographical comments, with a special reference to the Greek species. Pp. 287–295. In: Fet V. & Selden P., editors. *Scorpions 2001. In memoriam Gary A. Polis.* British Arachnological Society.
- Vachon, M. 1966. Liste des scorpions connus en Egypte, Arabie, Israel, Liban, Syrie, Jordanie, Turquie, Irak, Iran. *Toxicon*, 4:209–218.
- Vachon, M. 1949. Etudes sur les Scorpions. III (suite). Description des Scorpions du Nord de l'Afrique. *Archives de l'Institut Pasteur d'Algérie*, 27:66–100.
- Vachon, M. 1979. Etude des caractères utilisés pour classer les familles et les genres de Scorpions (Arachnides). *Bulletin du Muséum d'histoire Naturelle, Ser. Zoologie*, 3 (140):857–895.
- Wahbeh, Y. 1976 A study of Jordanian scorpions. *Jordan Medical Journal*, 11:84–92.